



Biscogniauxia (Hypoxylon) Canker and Dieback of Trees

Jennifer Olson
Plant Disease Diagnostician

Damon Smith
Extension Plant Pathology Specialist

Oklahoma Cooperative Extension Fact Sheets
are also available on our website at:
<http://osufacts.okstate.edu>

One of the greatest losses from the forested areas and landscapes in Oklahoma is the loss of a tree. Not only is a tree an aesthetic part of the landscape, but it also provides shade for humans as well as habitat for birds and other animals. In the landscape, trees can be the focal points for the entire planting. Therefore, it is tragic that a disease like Biscogniauxia (formerly Hypoxylon) canker and dieback can kill seemingly healthy trees in a very short period of time.

Occurrence

Although there were early records of Biscogniauxia canker and dieback on oaks (*Quercus* spp.) in Oklahoma, not much attention was given to its potential as an important disease of shade trees until 1979. Reports on the severity of this disease were first received from Pushmataha County in southeastern Oklahoma. A survey of several counties, and specimens received in the Plant Disease Diagnostic Laboratory, indicated that the disease was very active in at least 14 counties in the eastern and central portions of the state. It is now common throughout much of Oklahoma. Although all oak species are susceptible, the disease is more frequently observed on the red oak subgroup. While occurrence in Oklahoma is usually documented on oaks, other species of shade trees can be affected. Biscogniauxia canker and dieback can be found on maple (*Acer* spp.), hickory (*Carya* spp.), pecan (*Carya illinoensis*) and sycamore (*Platanus* spp.) among other hardwoods.

The Biscogniauxia canker and dieback fungus is found throughout the U.S. It has been diagnosed from several habitats, including forest sites; trees in pastures, recently developed home sites and established residential areas. In one case in Oklahoma, the severity of the disease eliminated 15 acres of established oak trees surrounding a recently built home. In another case, it eliminated six acres of oaks in the central area of a condominium development.

Disease Cycle

In Oklahoma, Biscogniauxia canker and dieback is reportedly caused by *Biscogniauxia atropunctata* (formerly *Hypoxylon atropunctatum*) and other *Biscogniauxia* spp. Biscogniauxia canker and dieback is a disease of stressed trees. The fungus colonizes healthy trees and lives in the bark and sapwood for an indefinite period of time without causing damage. In healthy trees, natural defenses are typically present that prevent the fungus from causing disease. When the tree becomes stressed, its defenses are reduced and the tree can no longer resist the fungus. Stress factors include drought, heat, root

damage, chemical injury and other diseases. Trees that are attacked by defoliating insects are also more likely to develop Biscogniauxia dieback. The most common stress factor in Oklahoma is drought and heat. The causal fungi are favored by warm and dry conditions. In fact, *B. atropunctata* can grow at temperatures up to 104 F and grows optimally at 95 F.

The fungus invades stressed trees through wounds. Early symptoms may appear as yellowing and wilting of leaves. The leaves will turn brown and the tops of branches are killed during the dieback phase (Figure 1) of the disease. This phase may go unnoticed or may be attributed to other problems. The disease will progress down to the trunk and cause a girdling canker, which kills the entire crown. At this point of the disease cycle, the entire tree will turn brown seemingly overnight (although girdling of the trunk actually takes several weeks to months). Internally, the fungus causes a brown discoloration of the sapwood. Subsequently, a yellow decay with black zone lines will develop.

When the diseased branches or trunk die, the fungus will form a cushion-like mat under the bark called a stroma (Figure 2). As the stroma develops, it exerts pressure between the wood and bark of the tree. Eventually, the pressure causes the bark to slough off the tree, revealing the stroma. At first, the stroma appears white to brown (Figure 3) and is covered by dusty asexual spores (conidia) of the fungus, which are easily blown to nearby trees. Later, the stroma becomes gray to black in color as the structures of the sexual stage are produced (Figure 4). An older stroma may be completely black in color. Spores are wind-blown, rain-splashed or carried by animals to nearby trees.

Control

There are no chemicals registered to treat Biscogniauxia canker and dieback of trees. In a commercial operation, where trees can be harvested for wood or pulp, it is recommended that trees be cut before decay reduces their value. In a home setting, individual trees that have more than 15 percent of the crown area infected should be cut to ground level and burned. Trees with less damage should be given extra care, such as watering during periods of drought, providing adequate fertilization and preventing damage to trunks or roots.

The fungus does remain active on dead wood, producing numerous spores. It is recommended to destroy diseased wood as soon as possible to prevent spore production and possible spread to nearby trees. This lessens the likelihood that nearby trees will become infected, however these trees still may become infected by wind-blown spores from other sources.



Figure 1. Canopy dieback in a mature oak tree as a result of *Biscogniauxia* canker and dieback



Figure 2. An exposed stroma caused by a species of *Biscogniauxia* on a scaffold limb of an oak tree

Prevention

There is no cure for *Biscogniauxia* canker and dieback. Therefore the best defense is to maintain trees in a healthy and vigorous growing condition by adequate fertilization and by providing adequate water throughout the year. Avoid wounding trees and prune out dead wood and declining limbs as they appear.



Figure 3. An exposed stroma caused by a species of *Biscogniauxia*. Note the presence of bark around the base of the tree as a result of the formation of the stroma.



Figure 4. Small, black, mature stroma caused by a species of *Biscogniauxia* on pecan (*Carya illinoensis*).

Oklahoma State University, in compliance with Title VI and VII of the Civil Rights Act of 1964, Executive Order 11246 as amended, Title IX of the Education Amendments of 1972, Americans with Disabilities Act of 1990, and other federal laws and regulations, does not discriminate on the basis of race, color, national origin, gender, age, religion, disability, or status as a veteran in any of its policies, practices, or procedures. This includes but is not limited to admissions, employment, financial aid, and educational services.

Issued in furtherance of Cooperative Extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, Director of Cooperative Extension Service, Oklahoma State University, Stillwater, Oklahoma. This publication is printed and issued by Oklahoma State University as authorized by the Vice President, Dean, and Director of the Division of Agricultural Sciences and Natural Resources and has been prepared and distributed at a cost of \$1.25 per copy. 0512 GH